



Extended IR Emission from (U)LIRGs

Vassilis Charmandaris
Univ. of Crete, Greece

Tanio Diaz-Santos (Univ. of Crete), Lee Armus (SSC/Caltech)
& the GOALS/IRS team

Papers:

Diaz-Santos et al. 2010 ApJ, 723, 993


Diaz-Santos et al. 2011 ApJ, (accepted - almost...)

Motivation

- A number of the observational properties of galaxies detected at sub-mm at $z \sim 2$ with $L_{\text{IR}} > 10^{12} L_{\odot}$ (SMG) such as:
 - *cold infrared colors*
 - *energy production dominated by star formation ($> 100 M_{\odot}/\text{yr}$)*
 - *mid-IR spectral features (ie PAH strength)*... resemble those of local LIRGs rather than ULIRGs.
- Kinematical evidence from ionized ($\text{H}\alpha$) and molecular (CO) gas are often consistent the presence of extended ($\sim 5\text{kpc}$) star forming disks
- There is a “broad” connection between mid-IR emission and star formation rate with some caveats: “main sequence/ compact-starburst” (Elbaz et al. 2011)
- We wish to quantify the extended extranuclear emission in local LIRGs in the $5\text{-}15\mu\text{m}$ range to contribute additional evidence in the analogy between the physics of the ISM excitation in LIRGs and the conditions seen in SMGs and ULIRGs at $z > 1$.

The Sample

- The sample is based on the Great Observatory Allsky LIRG Survey (GOALS; [Armus et al. 2009](#)) of 202 systems (181 of which are LIRGs)
- All systems are observed with all four Spitzer/IRS modules ($\sim 5\text{-}37\mu\text{m}$)
- Additional Spitzer data with IRAC/MIPS, as well as HST, GALEX, VLA, CO




The Method

- ❑ Use the 2D IRS longslit images of all sources and re-extract total 5-15 μ m spectrum using the same algorithms of the Spitzer pipeline
- ❑ Use a standard star (HR7341) as our unresolved point source (PSF)
- ❑ Scale the spatial profile of the standard star along the slit at every wavelength and subtract it from the corresponding profile of each source.
- ❑ Define as Fraction of Extended Emission (FEE):


$$\text{Fraction of EE } (\lambda) = \frac{\text{Total (U)LIRG flux } (\lambda) - \text{PSF } (\lambda)}{\text{Total (U)LIRG flux } (\lambda)}$$

Types of mid-IR spatial profiles




- Three spatial profiles are visually identified:
 - Constant:** no variation as a function of λ (~50% of sample),
 - PAH/line extended:** 20-70% of PAH flux is extended (~17% of sample),
 - Si "extended":** Si at 9.7 μm appears extended (~24% of sample) -> suggests that integrated spectrum underestimates nuclear extinction.

Extended Emission and L(IR)



The median fraction of extended emission decreases when L_{IR} increases.
Similarly for the 13.2 μm continuum emission

Extended Emission and Interaction stage




□ Use merger stage classification relying on optical/near-IR morphology from Petric et al. (2011 ApJ, 730,28)

□ 0: non interacting -> 5: mergers

□ More advanced mergers are more luminous and also more compact in their mid-IR continuum

(Similar to what has been shown in other wavelengths)

Extended Emission and AGN




□ Use mid-IR AGN classification (Petric et al. 2011) based on the “Laurent diagram” which probes the presence of a hot dust component (Laurent et al. 2000)

□ AGN dominated sources also more compact in their mid-IR continuum


Note that we refer to mid-IR dominant AGN, not bolometrically

Comparing the extent of various features




- Continuum at 13.2μm and 6.6μm as well as is [NeII] @12.8μm are equally extended.
- The 6.2μm and 7.7μm PAHs are as extended as the 13.2μm continuum
- The 11.3μm PAH is more extended as the 13.2μm continuum.
- Variation likely due to the ionized (6.2, 7.7μm) vs neutral (11.3μm) nature of PAHs. Ionized PAHs need harder radiation field to be excited ([Galliano et al 2008](#))

The spectrum of extended & compact components



- The 5-15 μm spectrum of the nuclear component varies depending on spectral type.
- The 5-15 μm spectrum of the extended component is similar for all three spectral types.
- Suggests common mechanism in the excitation (ie star formation) and dust properties
- Integrated spectra of SMGs (Menendez-Delmestre et al. 2009) display lower PAH EW, indicating the presence of a possible power-law hot dust continuum.

Extended Emission and far-IR Colors




- Use IRAS colors as a probe of the global ISM “temperature”
- Sources which are more compact in their mid-IR continuum have warmer far-IR colors
- This suggests that when nuclear emission is compact in the mid-IR it may dominate the energy production in the galaxy

Can be tested with Herschel Key Project: Hercules, Herschel-GOALS

$$FEE_{13.2\mu m} = 0.04 \pm 0.02 - (1.83 \pm 0.11) \times \log\left(\frac{f_{60\mu m}}{f_{100\mu m}}\right)$$

UV/mid-IR comparison of two LIRGs


Images: HST/STIS **UV** - Contours: ISO/CAM **7 μ m**



To be explored better with GOALS:

Preliminary GALEX/MIPS results (Howell et al. 2010) indicate that:

- Among interacting systems in GOALS 32% have one galaxy dominate the UV emission and the other the 24 μ m emission
- Based on number counts 15-30% of $z \sim 2$ systems are unresolved as VV114



7 μ m/UV ~ **800:10:35**

7 μ m/UV ~ **330:160:190**

The spatial resolution of ground & Spitzer/MIPS24 surveys of LIRGs at $z \sim 2$ will result in blending of the emission from the unresolved interacting components leading to a systematic underestimation of their dust content.

Conclusions / Perspectives

Even though the angular resolution of Spitzer/IRS is ~ 3.6 arcsecs..

- ❑ LIRGs display large fraction of extended mid-IR emission in both continuum and 5-15 μ m features
- ❑ For at least 90% of the sample more than 20% of the mid-IR flux originates outside the nuclear unresolved region.
- ❑ For at least 35% of the sample more than 50% of the mid-IR flux is extranuclear (probably also their star formation?)
- ❑ Systems with $\log(L_{\text{IR}}) > 11.8 L_{\odot}$ display mid-IR extent of less than 20%
- ❑ The 13.2 μ m size of LIRGs is ~ 3.5 kpc, while ULIRGs are less than 1.5 kpc
- ❑ The 11.3 μ m PAH emission is more extended, consistent with it being a "neutral" PAH.
- ❑ Spatial extent decrease with mid-IR AGN activity and merger stage.
- ❑ Compact sources have warmer far-IR colors

- ❑ To do:
Explore the implications of the mid-IR, UV, and radio continuum spatial extent of local (U)LIRGS in unresolved sources of deep high-z surveys